

Bijsturing van de respons bij stress-stoornissen



Roel de Rijk

DynaCorts Therapeutics BV


Department of Clinical Psychology

Prof.dr. W. van der Does

Department of Psychiatry

Prof.dr. B van Hemert

Antidepressiva gebruik in Nederland



De stijging van het antidepressivagebruik van 2011 tot 2012 is van dezelfde orde als het gemiddelde van alle geneesmiddelgroepen. Bron: Stichting Farmaceutische Kengetallen

Monoamine theory of depression

The Causes and Effects of Depression

Depression is a treatable illness caused by a decrease in neurotransmitter levels. Here's more information on the causes, effects, and treatment of depression.

SSRI's

TCA's

MAO-Inh



improvements



00:28

|| PAUSE

01:26



MENU

Stress en psychopathologie

Stress




CRH

AZS

HPA axis


Precipitatie psychopathologie




Is het mogelijk, met de huidige kennis van stress, een nieuw medicament te ontwikkelen

De snelle initiele stress-respons

Geanticipeerd
psychosociaal probleem



Corticosteroid receptor diversiteit: MR and GR in het brein



HPA as regulatie




Gedrags adaptatie

- 'Mineralocorticoid' Receptor*
- * hoge affiniteit
 - * discrete lokalisatie
 - * Hoge bezetting met cortisol

- Glucocorticoid Receptor*
- * 10x lagere affiniteit
 - * overall
 - * Bezet na stress

Reul/de Kloet 1985

Mineralocorticoid receptoren in de stress-respons en electrolyt regulatie




MR en perceptie: muizen

Ron de Kloet

Melly Oitzl

Gedrag's dynamiek

Test: zonder platform




Stress-respons maakt adaptieve actie mogelijk

Stressor

1. Evaluatie & emotie (MR)
2. Initiatie respons (CRH, NE)
3. Gedrag & fysiologie

4. Verhoogde niveaus cortisol
5. Activatie GR
6. Coordinatie van responsen



Hypothese:

Functionele genetische variantie van het MR-gen:

1. Determinant van de stress-respons
2. Kwestbaarheids factor


De “candidate gene approach”:

1. Breng humane genetische variatie in kaart
2. Karakteriseer genetische variatie in vitro
3. Specifieke in vivo testen

Approach:

1. Search for human MR-gene variants and haplotypes
2. Test functionality *in vitro*:
3. Cardiovascular regulation
4. Neuro-endocrine effects in healthy subjects
5. Behavior in healthy subjects
6. Association with psychopathology

MR-gene SNPs and haplotypes






Sequencing 50 anonymous subjects:

5' promotor, exons & 3' UTR regions



-2 G/C

I180V

Human MR-haplotypes are functional different



Verhoogde cortisol and autonome responsen tijdens de TSST in dragers van de MR180V




Preparation
speaking arithmetic


Public
Mental arithmetic

GR-SNPs are not confounding MR effects

MR I180V is associated with feelings of depression in the elderly



Gender dependent association between MR haplotype 2 and optimism




Association in elderly cohort from Arnhem

Overall mean optimism

Females: 1.35



Males: 1.34



LEIDS-R subscales in female students

questionnaire cognitive reactivity



Acceptation / coping



Aggression





Hopelessness



Perfectionism

Risk aversion







Rumination



Totaal (n= 140)

LEIDS-R subscales in female students




MR-gene variants → behavioral endophenotype / depression

Effect van genetische variatie in MR

	MR-variant	
	Hap 1&3	Hap 2
Optimisme	↓	↑
Hopeloosheid	↑	↓
Risco mijden	↑	↓
Piekeren	↑	↓
Aggressie	↑	↓
Neuroticisme	↑	↓
Depressie	↑	↓
	MR	MR

MR Hap 2 associates with less depression



□ Healthy controls
■ Patients

MR expression in depression?

Hypothesis: lower MR in depression


Q-PCR study, postmortem brain tissue

mRNA of MR, GR, exon1 α MR and exon1 β MR


MR-expression: controls vs. depressed

MR β role in neuronal protection


Hippocampus




Inferior frontal gyrus




Cingulate gyrus




Amygdala




Nucleus accumbens




Implications of human MR-gene variants on endophenotypes / depression



Fludrocortisone improves treatment of depression




PATIENTS COTREATED WITH SPIRONOLACTONE FOR 10 DAYS RESPONDED LESS FAVORABLY TO AMITRIPTYLINE THAN PATIENTS COTREATED WITH PLACEBO



MR als nieuwe behandeling van depressie

MR = cortisol / stress receptor



Personalized medicine: effect of functional genotypes

	<u>Dose agonist</u>		
<i>MR- genotype</i>	<u>low</u>	<u>middle</u>	<u>high</u>
<i>1/1; 3/3; 1/3</i>			X
<i>2/2</i>	X		
<i>1/2; 2/3</i>		X	

Nieuwe behandeling depressie: focus op stress-systeem



Selection of patients

Genotyping
Stress-measures
Others ?



MR-stimulation as add on
to SSRI's or psychotherapy



Improved and faster recovery

